

I-3-C Plus

Indole-3-Carbinol and DIM Supports Estrogen Metabolism*

I-3-C Plus is a potent blend of naturally occurring compounds that support healthy estrogen metabolism and promote healthy hormonal balance.⁺ The new and upgraded I-3-C Plus formula contains indole-3-carbinol (I-3-C) and diindolyImethane (DIM), both abundantly found in cruciferous vegetables. What makes the I-3-C Plus formula new and unique is that it now utilizes the clinically proven absorption of microencapsulated BioResponse DIM[®] Complex.⁺

How I-3-C Plus Works

The BioResponse DIM[®] Complex in the new I-3-C Plus is the only bioavailable, microencapsulated DIM formula for extended release and increased absorption. In its crystalline form, DIM is poorly absorbed. With patented BioResponse DIM[®] Complex, there is a statistically significant advantage of gastrointestinal absorption with microencapsulated DIM over the more commonly found crystalline DIM as shown in these graphical representations of results from clinical trials.^{+1,2,3,4}





Clinical research has shown that both I-3-C and DIM act on enzymes in the body that mediate estrogen production for both women and men.^{+5,6} I-3-C comes from the breakdown of a compound found in cruciferous vegetables called glucobrassicin—a type of glucosinolate. When ingested, the body breaks down I-3-C to DIM,



Form: 60 Capsules

Serving Size: 1 Capsule

Ingredients	Amount	% DV
Indole-3-Carbinol	150 mg	*
BioResponse DIM [®] Complex [starch, diindolylmethane (DIM), vitamin E (as tocophersolan), phosphatidyl choline, silica.]	50 mg	*

Other Ingredients:

Hypromellose, microcrystalline cellulose, tricalcium phosphate, vegetable magnesium stearate, silica.

BioResponse DIM[®] is a proprietary, enhanced bioavailability complex containing diindolylmethane licensed from BioResponse, L.L.C., Boulder, Colorado.

Directions:

Take one capsule up to three times daily as a dietary supplement, or as directed by your healthcare practitioner.

Caution: If you are pregnant, nursing, or taking medication, consult your healthcare practitioner before use. Keep out of reach of children.



These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.



ND239 Rev. 3/22 ©2022 NutriDyn

For more information, visit: www.nutridyn.com

both of which influence estrogen metabolism by acting on certain enzymes. In particular, I-3-C and DIM work by inhibiting the actions of histone deacetylase enzymes and inducing cytochrome P450 enzymes.^{5,6}

Both of these enzyme families are crucial for proper estrogen balance in the body. Specifically, I-3-C has been shown to induce an enzyme called CYP1A1, which aids in the metabolism of xenobiotics that can disrupt estrogen balance in the body.⁷ Absorbable BioResponse DIM[®] Complex works to activate histone deacetylase inhibitors, which research suggests modifies estrogen receptor binding to promote healthy estrogen metabolism and hormonal balance.^{+7,8,9,10}

Given this, both I-3-C and DIM appear to have significant effects on supporting healthy estrogen metabolism and promoting healthy hormonal balance.⁺ Clinical research also shows the proven bioavailability of BioResponse DIM[®] Complex to optimize the ratio of estrogen metabolites to support breast, uterine, cervical, and prostrate health.^{+1,2,3,4}

I-3-C Plus Supplementation

The ingredients in I-3-C Plus are dosed in a manner that is congruous with what research suggests to be effective and safe, particularly for supporting healthy estrogen metabolism and promoting healthy hormonal balance.⁺

Clinical evidence and research cited herein shows that the ingredients in I-3-C Plus may:

- Support healthy estrogen metabolism⁺
- Promote healthy hormonal balance⁺
- Promote balanced mood and well-being⁺

References:

- 1. Zeligs, M. A., Brownston, P. K., Sharp, M. E., Westerling, K. C., Wilson, S. M., & Johs, S. M. (2005). Managing cyclical mastalgia with absorbable diindolylmethane: A randomized, placebo-controlled trial. *Journal of the American Nutraceutical Association*, 8(1), 5-15.
- Zeligs, M. A., Fulfs, J. C., Peterson, R., Wilson, S. M., McIntyre, L., Sepkovic, D. W., & Bradlow, H. L. (2003). In vivo, uterine-protective activity of absorption-enhanced diindolylmethane: Animal and preliminary human use in combination with Tamoxi-fen. *Proceedings of the American* Association of Cancer Research, 44, 1268.
- **3.** Zeligs, M. A., Sepkovic, D. W., Manrique, C., Macsalka, M., Williams, D. E., & Bradlow, H. L. (2002). Absorption-enhanced 3,3'-diindolylmethane: Human use in HPV-related, benign and pre-cancerous conditions. *Proceedings of the American Association of Cancer Research, 43*, 3198.
- Teas, J., Cunningham, J. E., Fowke, J. H., Nitcheva, D., Kanwat, C. P., Boulware, R. J., Sepkovic, D. W., Hurley, T. G., & Hebert, J. R. (2005). Urinary estrogen metabolites, prostate specific antigen, and body mass index among African-American men in South Carolina. *Cancer Detection and Prevention*, 29(6), 494-500.
- 5. Michnovicz, J. J., Adlercreutz, H., & Bradlow, H. L. (1997). Changes in levels of urinary estrogen metabolites after oral indole-3-carbinol treatment in humans. *Journal of the National Cancer Institute*, 89(10), 718-723.
- Chen, I., McDougal, A., Wang, F., & Safe, S. (1998). Aryl hydrocarbon receptor-mediated antiestrogenic and antitumorigenic activity of diindolylmethane. *Carcinogenesis*, 19(9), 1631-1639.
- 7. Aggarwal, B. B., & Ichikawa, H. (2005). Molecular targets and anticancer potential of indole-3-carbinol and its derivatives. Cell Cycle, 4(9), 1201-1215.
- Leong, H., Riby, J. E., Firestone, G. L., & Bjeldanes, L. F. (2004). Potent ligand-independent estrogen receptor activation by 3,3'-diindolylmethane is mediated by cross talk between the protein kinase A and mitogen-activated protein kinase signaling pathways. *Molecular Endocrinology*, 18(2), 291-302.
- Dalessandri, K. M., Firestone, G. L., Fitch, M. D., Bradlow, H. L., & Bjeldanes, L. F. (2004). Pilot study: Effect of 3,3'-diindolylmethane supplements on urinary hormone metabolites in postmenopausal women with a history of early-stage breast cancer. *Nutrition and Cancer*, 50(2), 161-167.
- 10. Bradlow, H. L., Telang, N. T., Sepkovic, D. W., & Osborne, M. P. (1996). 2-hydroxyestrone: the 'good' estrogen. *Journal of Endocrinology*, 150(S2), 59-65.

For more information, visit: www.nutridyn.com